

§1.2: IDEALS

DEFINITION: Let S be a subset of a ring R . The **ideal generated by S** , denoted (S) , is the smallest ideal containing S . Equivalently,

$$(S) = \left\{ \sum r_i s_i \mid r_i \in R, s_i \in S \right\} \quad \text{is the set of } R\text{-linear combinations}^1 \text{ of elements of } S.$$

We say that S **generates** an ideal I if $(S) = I$.

DEFINITION: Let I, J be ideals of a ring R . The following are ideals:

- $IJ := (ab \mid a \in I, b \in J)$.
- $I^n := \underbrace{I \cdot I \cdots I}_{n \text{ times}} = (a_1 \cdots a_n \mid a_i \in I)$ for $n \geq 1$.
- $I + J := \{a + b \mid a \in I, b \in J\} = (I \cup J)$.
- $rI := (r)I = \{ra \mid a \in I\}$ for $r \in R$.
- $I : J := \{r \in R \mid rJ \subseteq I\}$.

DEFINITION: Let I be an ideal in a ring R . The **radical** of I is $\sqrt{I} := \{f \in R \mid f^n \in I \text{ for some } n \geq 1\}$. An ideal I is **radical** if $I = \sqrt{I}$.

DIVISION ALGORITHM: Let A be a ring, and $R = A[X]$ be a polynomial ring. Let $g \in R$ be a **monic** polynomial; i.e., the leading coefficient of f is a unit. Then for any $f \in R$, there exist unique polynomials $q, r \in R$ such that $f = gq + r$ and the top degree of r is less than the top degree of g .

(1) Briefly discuss why the two characterizations of (S) in Definition 2.1 are equal.

The set of linear combinations of elements of S is an ideal:

- $0 = 0s_1$ (we also consider 0 to be the empty combination);
- given two linear combinations, by including zero coefficients, we can assume our combinations involve the same elements of S , and then $\sum_i a_i s_i + \sum_i b_i s_i = \sum_i (a_i + b_i) s_i$;
- $r(\sum_i a_i s_i) = \sum_i r a_i s_i$.

Any ideal that contains S must contain all of the linear combinations of S , using the definition of ideal. These two facts mean that the set of linear combinations is the smallest ideal containing S .

(2) Finding generating sets for ideals: Let S be a subset of a ring R , and I an ideal.

- (a)** To show that $(S) = I$, which containment do you think is easier to verify? How would you check?
- (b)** To show that $(S) = I$ given $(S) \subseteq I$, explain why it suffices to show that $I/(S) = 0$ in $R/(S)$; i.e., that every element of I is equivalent to 0 modulo S .
- (c)** Let K be a field, $R = K[U, V, W]$ and $S = K[X, Y]$ be polynomial rings. Let $\phi : R \rightarrow S$ be the ring homomorphism that is constant on K , and maps $U \mapsto X^2, V \mapsto XY, W \mapsto Y^2$. Show that the kernel ϕ is generated by $V^2 - UW$ as follows:
 - Show that $(V^2 - UW) \subseteq \ker(\phi)$.
 - Think of R as $K[U, W][V]$. Given $F \in \ker(\phi)$, use the Division Algorithm to show that $F \equiv F_1 V + F_0$ modulo $(V^2 - UW)$ for some $F_1, F_0 \in K[U, W]$ with $F_1 V + F_0 \in \ker(\phi)$.
 - Use $\phi(F_1 V + F_0) = 0$ to show that $F_1 = F_0 = 0$, and conclude that $F \in \ker(\phi)$.

(a) Showing $(S) \subseteq I$ is the easier containment: it suffices to show that $S \subseteq I$.

(b) This follows from the Second Isomorphism Theorem.

¹Linear combinations always means *finite* linear combinations: the axioms of a ring can only make sense of finite sums.

- (c)
- We check $\phi(V^2 - UW) = (XY)^2 - X^2Y^2 = 0$, so $V^2 - UW \in \ker(\phi)$. This implies $(V^2 - UW) \subseteq \ker(\phi)$.
 - By Division, we have $F = (V^2 - UW)Q + R$, with the top degree (in V) of R at most 1. Then $F \equiv R = F_1V + F_0$ modulo $(V^2 - UW)$. Since $F, V^2 - UW \in \ker(\phi)$, we must have $F_1V + F_0 \in \ker(\phi)$.
 - We have $0 = \phi(F_1V + F_0) = F_1(X^2, Y^2)XY + F_0(X^2, Y^2)$. The $F_1(X^2, Y^2)XY$ terms only have monomials whose X -degree is odd, and the $F_0(X^2, Y^2)$ terms only have monomials whose X -degree is even, so none can cancel with each other. This means that $F_1(X^2, Y^2) = 0$ and $F_0(X^2, Y^2) = 0$, so $F_1(U, W) = F_0(U, W) = 0$. Thus, $F \equiv 0$ modulo $(V^2 - UW)$, and as above, we conclude $\ker(\phi) = (V^2 - UW)$.

(3) Radical ideals:

(a) Fill in the blanks and convince yourself:

- R/I is a field $\iff I$ is _____
- R/I is a domain $\iff I$ is _____
- R/I is reduced $\iff I$ is _____

(b) Show that the radical of an ideal is an ideal.

(c) Show that a prime ideal is radical.

(d) Let K be a field and $R = K[X, Y, Z]$. Find a generating set² for $\sqrt{(X^2, XYZ, Y^2)}$.

(a)

- R/I is a field $\iff I$ is maximal
- R/I is a domain $\iff I$ is prime
- R/I is reduced $\iff I$ is radical

(b) Let $f, g \in \sqrt{I}$. Then there are $m, n \geq 1$ such that $f^m, g^n \in I$. Then

$$(f + g)^{m+n-1} = \sum_{i+j=m+n-1} \binom{m+n-1}{i, j} f^i g^j,$$

and for each term in the sum either $i \geq m$ or $j \geq n$, so each term is in I , hence the whole sum is in I . Now let $r \in R$. Then $(rf)^m = r^m f^m \in I$.

(c) Suppose I is prime. If $x \in \sqrt{I}$, then $x^n \in I$ for some n . Then, by the definition of prime, $x \in I$. Thus, $\sqrt{I} = I$.

(d) Since X^2 and Y^2 are in (X^2, XYZ, Y^2) , we have $X, Y \in \sqrt{(X^2, XYZ, Y^2)}$ by definition, so $(X, Y) \subseteq \sqrt{(X^2, XYZ, Y^2)}$. For the other containment, if $F(X, Y, Z) \notin (X, Y)$, consider F as a polynomial in X, Y with coefficients in $K[Z]$; the condition means that the top degree of F is zero, and hence the top degree of F^n is zero for all n , so $F \notin \sqrt{(X^2, XYZ, Y^2)}$.

(4) Evaluation ideals in polynomial rings: Let K be a field and $R = K[X_1, \dots, X_n]$ be a polynomial ring. Let $\alpha = (\alpha_1, \dots, \alpha_n) \in K^n$.

(a) Let $\text{ev}_\alpha : R \rightarrow K$ be the map of evaluation at α : $\text{ev}_\alpha(f) = f(\alpha_1, \dots, \alpha_n)$, or $f(\alpha)$ for short. Show that $\mathfrak{m}_\alpha := \ker \text{ev}_\alpha$ is a maximal ideal and $R/\mathfrak{m}_\alpha \cong K$.

(b) Apply division repeatedly to show that $\mathfrak{m}_\alpha = (X_1 - \alpha_1, \dots, X_n - \alpha_n)$.

(c) For $K = \mathbb{R}$ and $n = 1$, find a maximal ideal that is not of this form. Same question with $n = 2$.

(d) With K arbitrary again, show that every maximal ideal \mathfrak{m} of R for which $R/\mathfrak{m} \cong K$ is of the form \mathfrak{m}_α for some $\alpha \in K^n$. Note: this is *not* a theorem with a fancy German name.

²Hint: To show your set generates, you might consider the bottom degree of F considered as a polynomial in X and Y .

- (a) The evaluation map is surjective, since for any $k \in K$, the constant function k maps to k . By the First Isomorphism Theorem, $R/\mathfrak{m}_\alpha \cong K$, so \mathfrak{m}_α is maximal.
- (b) We have $\text{ev}_\alpha(X_i - \alpha_i) = \alpha_i - \alpha_i = 0$, so $(X_1 - \alpha_1, \dots, X_n - \alpha_n) \subseteq \mathfrak{m}_\alpha$. Given some $F \in \mathfrak{m}_\alpha$, consider F as a polynomial in X_1 and apply division by $X_1 - \alpha_1$, to get $F \equiv F_1$ modulo $(X_1 - \alpha_1, \dots, X_n - \alpha_n)$, for some F_1 not involving X_1 . Continue with $X_2 - \alpha_2, \dots$ to get the F is equivalent to a constant, which must be zero. This shows that $F \in (X_1 - \alpha_1, \dots, X_n - \alpha_n)$, so $\mathfrak{m}_\alpha = (X_1 - \alpha_1, \dots, X_n - \alpha_n)$.
- (c) $(X^2 + 1); (X^2 + 1, Y)$.
- (d) Let $\phi : R \rightarrow R/\mathfrak{m} \cong K$ be quotient map followed by the given isomorphism. Set $\alpha_i := \phi(X_i)$. Then $X_i - \alpha_i \in \ker(\phi)$, so $\mathfrak{m}_\alpha = (X_1 - \alpha_1, \dots, X_n - \alpha_n) \subseteq \ker(\phi)$. Since \mathfrak{m}_α is maximal, we must have equality.

(5) Lots of generators:

- (a) Let K be a field and $R = K[X_1, X_2, \dots]$ be a polynomial ring in countably many variables. Explain³ why the ideal $\mathfrak{m} = (X_1, X_2, \dots)$ cannot be generated by a finite set.
- (b) Show that the ideal $(X^n, X^{n-1}Y, \dots, XY^{n-1}, Y^n) \subseteq K[X, Y]$ cannot be generated by fewer than $n + 1$ generators.
- (c) Let $R = \mathcal{C}([0, 1], \mathbb{R})$ and $\alpha \in (0, 1)$. Show that for any element $g \in (f_1, \dots, f_n) \subseteq \mathfrak{m}_\alpha$, there is some $\varepsilon > 0$ and some $C > 0$ such that $|g| < C \max_i\{|f_i|\}$ on $(\alpha - \varepsilon, \alpha + \varepsilon)$. Use this to show that \mathfrak{m}_α cannot be generated by a finite set.

- (a) Suppose $\mathfrak{m} = (f_1, \dots, f_m)$. Since each polynomial involves only finitely many variables, only finitely many variables occur in $\{f_1, \dots, f_m\}$, and since each f_i has no constant term, these polynomials are linear combinations of those variables X_1, \dots, X_n ; i.e., $(f_1, \dots, f_m) \subseteq (X_1, \dots, X_n)$. It suffices to show that $\mathfrak{m} \neq (X_1, \dots, X_n)$. To see it, take X_{n+1} and note that $X_{n+1} = \sum_{i=1}^n g_i X_i$ is impossible, since the monomial X_{n+1} can't occur in any summand of the right hand side.
- (b) Note that this ideal is the set of all polynomial whose bottom degree is at least n . Given a generating set f_1, \dots, f_m for I , consider the degree n terms of the polynomials f_i . We claim that the degree n terms of f_1, \dots, f_m must span the space of degree n polynomials as a vector space. Indeed, given h of degree n , we have $h \in I$, so $h = \sum_i g_i f_i$. But every term of f_i has degree at least n , so the only things of degree n on the right hand side come from the degree n piece of f_i and the degree zero piece of g_i . This shows the claim. Then the statement is clear, since the degree n terms form an $n + 1$ dimensional vector space.
- (c) Let $g = \sum g_i f_i \in (f_1, \dots, f_n)$. By continuity, there is some $\varepsilon > 0$ and some $C > 0$ such that $|g_i| < C/n$ on $(\alpha - \varepsilon, \alpha + \varepsilon)$, so $|g| < |\sum_i g_i f_i| \leq \sum_i |g_i| |f_i| \leq \sum_i C/n \max_i\{|f_i|\} \leq C \max_i\{|f_i|\}$ on $(\alpha - \varepsilon, \alpha + \varepsilon)$.
Now, given $f_1, \dots, f_n \in \mathfrak{m}_\alpha$, let $g = \sqrt{\max_i\{|f_i|\}}$. Then g is continuous and $g(\alpha) = 0$, so $g \in \mathfrak{m}_\alpha$, but $g/\max_i\{|f_i|\} = 1/g \rightarrow \infty$ as $x \rightarrow \alpha$, so there is no constant $C > 0$ and no interval $(\alpha - \varepsilon, \alpha + \varepsilon)$ on which $|g| < C \max_i\{|f_i|\}$. Thus, \mathfrak{m}_α is not finitely generated.

(6) Evaluation ideals in function rings: Let $R = \mathcal{C}([0, 1], \mathbb{R})$. Let $\alpha \in [0, 1]$.

- (a) Let $\text{ev}_\alpha : \mathcal{C}([0, 1]) \rightarrow \mathbb{R}$ be the map of evaluation at α : $\text{ev}_\alpha(f) = f(\alpha)$. Show that $\mathfrak{m}_\alpha := \ker \text{ev}_\alpha$ is a maximal ideal and $R/\mathfrak{m}_\alpha \cong \mathbb{R}$.
- (b) Show that $(x - \alpha) \subseteq \mathfrak{m}_\alpha$.

³Hint: You might find it convenient to show that $(f_1, \dots, f_m) \subseteq (X_1, \dots, X_n)$ for some n , and then show that $(X_1, \dots, X_n) \subsetneq \mathfrak{m}$

- (c) Show that every maximal ideal R is of the form \mathfrak{m}_α for some $\alpha \in [0, 1]$. You may want to argue by contradiction: if not, there is an ideal I such that the sets $U_f := \{x \in [0, 1] \mid f(x) \neq 0\}$ for $f \in I$ form an open cover of $[0, 1]$. Take a finite subcover U_{f_1}, \dots, U_{f_t} and consider $f_1^2 + \dots + f_t^2$.

- (a) $\text{ev}_\alpha : \mathcal{C}([0, 1]) \rightarrow \mathbb{R}$ is a surjective ring homomorphism, since $\text{ev}_\alpha(r) = r$ for any $r \in \mathbb{R}$. Thus, by the First Isomorphism Theorem, $R/\mathfrak{m}_\alpha \cong \mathbb{R}$, and hence \mathfrak{m}_α is a maximal ideal.
- (b) It suffices to note that $\text{ev}_\alpha(x - \alpha) = 0$.
- (c) Argue by contradiction: if not, there is a proper ideal I that is not contained in some \mathfrak{m}_α ; this means that for every α , some element of I does not vanish at α . Since for any continuous f , the set $U_f := \{x \in [0, 1] \mid f(x) \neq 0\}$ is open, the collection $\{U_f \mid f \in I\}$ is an open cover of $[0, 1]$. Since $[0, 1]$ is compact, there is a finite subcover U_{f_1}, \dots, U_{f_t} . For these f_i 's consider $h = f_1^2 + \dots + f_t^2$. Each f_i^2 is nonnegative, and for any α , one of these is strictly positive at α . This means that $h(x) \neq 0$ for all $x \in [0, 1]$, so h is a unit, and hence $I = R$, a contradiction.

(7) Division Algorithm.

- (a) What fails in the Division Algorithm when g is not monic? Uniqueness? Existence? Both?
- (b) Review the proof of the Division Algorithm.

- (8) Let K be a field and $R = K[[X_1, \dots, X_n]]$ be a power series ring in n indeterminates. Let $R' = K[[X_1, \dots, X_{n-1}]]$, so we can also think of $R = R'[[X_n]]$. In this problem we will prove the useful analogue of division in power series rings:

WEIERSTRASS DIVISION THEOREM: Let $r \in R$, and write $g = \sum_{i \geq 0} a_i X_n^i$ with $a_i \in R'$. For some $d \geq 0$, suppose that $a_d \in R'$ is a unit, and that $a_i \in R'$ is *not* a unit for all $i < d$. Then, for any $f \in R$, there exist unique $q \in R$ and $r \in R'[X_n]$ such that $f = qg + r$ and the top degree of r as a polynomial in X_n is less than d .

- (a) Show the theorem in the very special case $g = X_n^d$.
- (b) Show the theorem in the special case $a_i = 0$ for all $i < d$.
- (c) Show the uniqueness part of the theorem.⁴
- (d) Show the existence part of the theorem.⁵

- (a) Given f , write $f = \sum_{i \geq 0} b_i X_n^i$ with $b_i \in R'$. For existence, just take $r = \sum_{i=0}^{d-1} b_i X_n^i$ and $q = \sum_{i=d}^{\infty} b_i X_n^{i-d}$. For uniqueness, note that if $f = qg + r = q'g + r'$ with the top degree of r and r' as polynomials in X_n are less than d . Then $0 = g(q - q') + (r - r')$, so the uniqueness claim reduces to the case $f = 0$; we will use this in the other parts without comment. Every term of r has X_n -degree less than d , whereas every term of qg has X_n -degree at least d , so no terms can cancel. Thus $qg + r = 0$ implies $q = r = 0$ (here and henceforth, we assume r is as in the statement when we write $qg + r$).
- (b) If $a_i = 0$ for $i < d$, then $g = X_n^d u$ where $u = \sum_{i \geq 0} a_{i+d} X_n^i$. Since the constant coefficient of u is a_d , which is a unit in R' , u is a unit in R . Thus, we can apply (a) to f and X_n^d to get

⁴Hint: For an element of R' or of R , write ord' for the order in the X_1, \dots, X_{n-1} variables; that is, the lowest total X_1, \dots, X_{n-1} -degree of a nonzero term (not counting X_n in the degree). If $qg + r = 0$, write $q = \sum_i b_i X_n^i$. You might find it convenient to pick i such that $\text{ord}'(b_i)$ is minimal, and in case of a tie, choose the smallest such i among these.

⁵Hint: Write $g_- = \sum_{i=0}^{d-1} a_i X_n^i$ and $g_+ = \sum_{i=d}^{\infty} a_i X_n^i$. Apply (b) with g_+ instead of g , to get some q_0, r_0 ; write $f_1 = f - (q_0 g_+ + r_0)$, and keep repeating to get a sequence of q_i 's and r_i 's. Show that $\text{ord}'(q_i), \text{ord}'(r_i) \geq i$, and use this to make sense of $q = \sum_i q_i$ and $r = \sum_i r_i$.

$f = q_0X_n^d + r_0 = (q_0u^{-1})g + r_0$; thus, $q = q_0u^{-1}$ and $r = r_0$ satisfy the existence clause of the theorem. For uniqueness, if $f = q'g + r'$, then $f = q'uX_n^d + r'$, so by the uniqueness part of (a), we must have $q'u = q_0$ and $r' = r_0$, and thus $q' = q$ and $r' = r$.

(c) For an element of R' or of R , write ord' for the order in the X_1, \dots, X_{n-1} variables; that is, the lowest total X_1, \dots, X_{n-1} -degree of a nonzero term (not counting X_n in the degree). Suppose that $qg + r = 0$, and write $q = \sum_i b_i X_n^i$. Suppose that q is nonzero, so $b_i \neq 0$ for some i . Pick i such that $\text{ord}'(b_i) \leq \text{ord}'(b_j)$ for all j with $b_j \neq 0$, and $\text{ord}'(b_i) = \text{ord}'(b_j)$ implies $i < j$; we can do this by well ordering of \mathbb{N} . Say $\text{ord}'(b_i) = t$. Consider the coefficient of X_n^{d+i} in $0 = qg + r$. By the degree constraint on r , this is the same as the coefficient of X_n^{d+i} in qg . Multiplying out, this is $\sum_{j=0}^{d+i} a_{d+i-j} b_j$. For $j = i$, the order of $a_d b_i$ is t . For $j < i$, we have $\text{ord}'(a_{d+i-j} b_j) \geq \text{ord}'(b_j) > t$ by choice of i . For $j > i$, since $\text{ord}'(a_{d+i-j}) > 0$ and $\text{ord}'(b_j) \geq t$, we have $\text{ord}'(a_{d+i-j} b_j) > t$. Thus, the no term can cancel the $a_d b_i$ term, so $qg + r \neq 0$. On the other hand, if $q = 0$ and $r \neq 0$, clearly $qg + r \neq 0$. It follows there there are unique q, r such that $qg + r = 0$.

(d) First, we observe that in the context of (b), if $\text{ord}'(f) = t$, then $\text{ord}'(q), \text{ord}'(r) \geq t$. This is clear in the setting of (a), and following the proof of (b), we just need to observe that if u is a unit in R , then $\text{ord}'(q_0u^{-1}) \geq \text{ord}'(q_0)$, which is clear since any coefficient of the product q_0u^{-1} is a sum of multiples of the coefficients of q_0 .

Now we begin the main proof. Write $g_- = \sum_{i=0}^{t-1} a_i X_n^i$ and $g_+ = \sum_{i=t}^{\infty} a_i X_n^i$. Apply (b) with g_+ to write $f = q_0g_+ + r_0$, and set $f_1 = f - (q_0g_+ + r_0) = -q_0g_-$. Repeat with f_1 to write $f_1 = q_1g_+ + r_1$, and $f_2 = f_1 - (q_1g_+ + r_1) = -q_1g_-$. Continue like so to obtain a sequence of series q_0, q_1, \dots and r_0, r_1, \dots . From the observation above, we have that $\text{ord}'(q_i), \text{ord}'(r_i) \geq \text{ord}'(f_i) \geq \text{ord}'(q_{i-1}) + 1$, since the constant term of each coefficient of g_- vanishes. It follows that $\text{ord}'(q_i), \text{ord}'(r_i) \geq i$ for each i .

For a series h , write $[h]_i$ for the degree i part of h , and $[h]_{\leq i}$ for the sum of all parts of degree $\leq i$. Define q to be the series such that $[q]_i = \sum_{j=0}^i [q_j]_i$, and likewise with r . Note that r is still a polynomial in X_n of top degree less than d . We claim that $f = qg + r$. To show this, it suffices to show that $[f]_i = [qg + r]_i$. Note that to compute $[qg + r]_i$, we can replace q, g, r by $[q]_{\leq i}$, and similarly for the others. But $[q]_{\leq i} = [\sum_{j=0}^i q_j]_{\leq i}$ (and likewise with r), so $[qg + r]_i = [(\sum_{j=0}^i q_j)g + (\sum_{j=0}^i r_j)]_i$. Then, by construction of the sequences $\{q_i\}, \{r_i\}, \{f_i\}$, we have $[f - (qg + r)]_i = [f_{i+1}]_i$ and since $\text{ord}'(f_{i+1}) \geq i + 1$, we have $[f_{i+1}]_i = 0$. It follows that $f - (qg + r) = 0$; i.e., $f = qg + r$.