

§1.5: DETERMINANTS

Recall that given matrices A and B , the matrix product AB consists of linear combinations, namely: Each column of AB is a linear combinations of the columns of A , with coefficients/weights coming from the corresponding columns of B . That is,

$$(\text{col } j \text{ of } AB) = \sum_{i=1}^t b_{ij} \cdot (\text{col } i \text{ of } A);$$

note that b_{1j}, \dots, b_{tj} is the j -th column of B .

PROPERTIES OF det: For a ring R , the determinant is a function $\det : \text{Mat}_{n \times n}(R) \rightarrow R$ such that:

- (1) \det is a polynomial expression of the entries of A of degree n .
- (2) \det is a linear function of each column.
- (3) $\det(A) = 0$ if the columns are linearly dependent.
- (4) $\det(AB) = \det(A) \det(B)$.
- (5) \det can be computed by Laplace expansion along a row/column.
- (6) $\det(A) = \det(A^{\text{tr}})$.
- (7) If $\phi : R \rightarrow S$ is a ring homomorphism, and $\phi(A)$ is the matrix obtained from A by applying ϕ to each entry, then $\det(\phi(A)) = \phi(\det(A))$.

ADJOINT TRICK: For an $n \times n$ matrix A over R ,

$$\det(A) \mathbb{1}_n = A^{\text{adj}} A = A A^{\text{adj}},$$

where $(A^{\text{adj}})_{ij} = (-1)^{i+j} \det(\text{matrix obtained from } A \text{ by removing row } j \text{ and column } i)$.

EIGENVECTOR TRICK: Let A be an $n \times n$ matrix, $v \in R^n$, and $r \in R$. If $Av = rv$, then $\det(r \mathbb{1}_n - A)v = 0$. Likewise, if instead v is a row vector and $vA = rv$, then $\det(r \mathbb{1}_n - A)v = 0$.

DEFINITION: Given an $n \times m$ matrix A and $1 \leq t \leq \min\{m, n\}$ the **ideal of $t \times t$ minors of A** , denoted $I_t(A)$, is the ideal generated by the determinants of all $t \times t$ submatrices of A given by choosing t rows and t columns. For $t = 0$, we set $I_0(A) = R$ and for $t > \min\{m, n\}$ we set $I_t(A) = 0$.

LEMMA: If A is an $n \times m$ matrix, B is an $m \times \ell$ matrix, and $t \leq 1$, then

- $I_{t+1}(A) \subseteq I_t(A)$
- $I_t(AB) \subseteq I_t(A) \cap I_t(B)$.

PROPOSITION: Let M be a finitely presented module. Suppose that A is an $n \times m$ presentation matrix for M . Then $I_n(A)M = 0$. Conversely, if $fM = 0$, then $f^n \in I_n(A)$.

- (1)** Let M be a module. Suppose that m_1, \dots, m_n is a generating set with corresponding presentation matrix A . Which of the following is true:

$$A \begin{bmatrix} m_1 \\ \vdots \\ m_n \end{bmatrix} \stackrel{?}{=} 0 \qquad [m_1 \ \cdots \ m_n] A \stackrel{?}{=} 0.$$

Explain your answer in terms of the recollection on matrix multiplication above.

- (2) Eigenvector Trick:
- (a) What familiar fact/facts from linear algebra (over fields) is/are related to the Eigenvector Trick?
 - (b) Use the Adjoint Trick to prove the Eigenvector Trick.
- (3) Show that a square matrix over a ring R is invertible if and only if its determinant is a unit.
- (4) Proof of Proposition:
- (a) First consider the case $m = n$. Show that $\det(A)$ kills each generator m_i , and conclude that $I_n(A)M = 0$.
 - (b) Now consider the case $n \leq m$. Show that for any $n \times n$ submatrix A' of A that $\det(A')M = 0$, and conclude that $I_n(A)M = 0$. What's the deal when $m < n$?
 - (c) For the “conversely” statement, show that if $fM = 0$ then there is some matrix B such that $AB = f\mathbb{1}_n$, and deduce that $f \in I_n(A)^n$.
- (5) Prove the Lemma above.
- (6) Prove¹ FITTING'S LEMMA: If A and B are presentation matrices for the same R -module M of size $n \times m$ and $n' \times m'$ (respectively), and $t \geq 0$, then $I_{n-t}(A) = I_{n'-t}(B)$.

¹Hint: First consider the case when the two presentations have the same generating sets, but different generating sets for the relations. Reduce to the case where $B = [A|v]$ for a single column v .