

IDEALS

DEFINITION: Let R be a ring. An **ideal** of R (also called a **two-sided ideal**) is a nonempty subset of R such that

- (1) I is closed under addition: for all $a, b \in I$, we have $a + b \in I$.
- (2) I absorbs multiplication: for all $r \in R$ and $a \in I$, we have $ra \in I$ and $ar \in I$.

A **left ideal** of R is a nonempty subset of R such that

- (1) I is closed under addition: for all $a, b \in I$, we have $a + b \in I$.
- (2ℓ) I absorbs left multiplication: for all $r \in R$ and $a \in I$, we have $ra \in I$.

The definition of **right ideal** is analogous

LEMMA 1 (GENERAL RECIPES FOR IDEALS): Let R be a ring.

- (i) If I, J are ideals, then $I + J := \{a + b \mid a \in I, b \in J\}$ is an ideal.
- (ii) If I, J are ideals, then $IJ := \{\sum_{i=1}^n a_i b_i \mid a_i \in I, b_i \in J\}$ is an ideal.
- (iii) If $\{I_\alpha\}_{\alpha \in A}$ is an arbitrary collection of ideals of R , then $\bigcap_{\alpha \in A} I_\alpha$ is an ideal.
- (iv) If $\{I_\alpha\}_{\alpha \in A}$ is a *chain*¹ of ideals, then $\bigcup_{\alpha \in A} I_\alpha$ is an ideal.
- (v) If I is an ideal, then $I[x] := \{a_0 + a_1 x + \cdots + a_n x^n \mid a_i \in I\}$ is an ideal of $R[x]$.

(1) Working with the definition:

- (a) If I is an ideal (or left ideal) of R , explain why $0 \in I$ and $(I, +)$ is a subgroup of $(R, +)$.
- (b) Very quickly explain why $\{0\}$ and R are ideals of R . We say that an ideal is **nontrivial** if $I \neq 0$ and proper if $I \neq R$.
- (c) Explain why an ideal $I \subseteq R$ is proper if and only if $1 \notin I$.
- (d) Quickly explain why “ideal,” “left ideal,” and “right ideal” are identical notions in a commutative ring.

(2) Show that the subset

$$\left\{ \begin{bmatrix} a & 0 \\ b & 0 \end{bmatrix} \mid a, b \in \mathbb{R} \right\} \subseteq \text{Mat}_2(\mathbb{R})$$

is a left ideal, but is not a (two-sided) ideal.

(3) Prove parts (i) and (v) of Lemma 1.

(4) Show² that the union of two ideals does not have to be an ideal in general.

(5) Prove parts (ii)–(iv) of Lemma 1.

¹This means that for all $\alpha, \beta \in A$, either $I_\alpha \subseteq I_\beta$ or $I_\beta \subseteq I_\alpha$.

²Hint: Consider $2\mathbb{Z}, 3\mathbb{Z} \subseteq \mathbb{Z}$.

DEFINITION: Let R be a ring. and $S \subseteq R$ be a subset. The **ideal generated by S** is the ideal

$$(S) = \bigcap_{\substack{I \text{ ideal} \\ I \supseteq S}} I.$$

An ideal is **principal** if $I = (a)$ for a single element $a \in R$.

LEMMA 2 (IDEAL GENERATED A SUBSET): Let R be a ring and $S \subseteq R$.

- (i) There is an equality $(S) = \{\sum_{i=1}^n r_i a_i r'_i \mid r_i, r'_i \in R, a_i \in S\}$.
- (ii) If R is commutative, then $(S) = \{\sum_{i=1}^n r_i a_i \mid r_i \in R, a_i \in S\}$.
- (iii) If R is commutative and $a \in R$, then $(a) = \{ra \mid r \in R\}$.

(6) Let $R = \mathbb{Z}[x]$. Use the Lemma to quickly explain the following:

- (a) (2) is the set of all integer polynomials with every coefficient even.
- (b) (x) is the set of all integer polynomials with zero constant term.
- (c) $(2, x)$ is the set of all integer polynomials with even constant term.

(7) Show³ that the ideal $(2, x)$ in $\mathbb{Z}[x]$ is not principal.

(8) Show that if R is noncommutative then one can⁴ have $(a) \supsetneq \{rar' \mid r, r' \in R\}$.

³Hint: If $(2, x) = (f)$, with $f = a_0 + a_1x + \cdots + a_nx^n$, note that $2, x \in (f)$. What can you say about a_0 ?

⁴Hint: You can use the fact that you will prove in HW#11 that $\text{Mat}_n(F)$ has no nontrivial proper ideals if F is a field.